Write your name here		0.1	
Surname		Other name	s
Pearson Edexcel Level 1/Level 2 GCSE (9-1)	Centre Number		Candidate Number
<b>Mathemat</b>	icc		)
	ICS.		
Paper 2 (Calculator)	.ICS		
	.ICS		Higher Tier
			Paper Reference
Paper 2 (Calculator)			

## **Instructions**

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
   there may be more space than you need.
- You must **show all your working**.
- Diagrams are **NOT** accurately drawn, unless otherwise indicated.
- Calculators may be used.
- If your calculator does not have a  $\pi$  button, take the value of  $\pi$  to be 3.142 unless the question instructs otherwise.

## Information

- The total mark for this paper is 80
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

## **Advice**

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



P48528A
©2018 Pearson Education Ltd.
6/7/7/7/8/7/



## Answer ALL questions.

Write your answers in the spaces provided.

You must write down all the stages in your working.

(a) Simplify  $m^3 \times m^4$ 

Laws of Indices

$$x_{\alpha} \times x_{p} = x_{\alpha+p}$$

$$M^{3+4} = m^7$$

$$m^7$$

(b) Simplify  $(5np^3)^3$ 

$$(5 \times 1 \times 1)^3 = \text{raise each term to the power of }$$

$$= 5^{3} \times n^{3} \times (p^{3})^{3} \qquad (x^{a})^{b} = x^{ab}$$

$$= 125 \times n^{3} \times p^{9} \qquad (p^{3})^{3} = p^{3 \times 3} = p^{9}$$

$$= 125 n^{3} p^{9} \qquad (p^{3})^{3} = p^{3 \times 3} = p^{9}$$

$$-\left(x^{\alpha}\right)^{b}=x^{\alpha b}$$

$$\left(b_{3}\right)_{3}=b_{3\times 9}=b_{0}$$

1) 2 correct tems

(c) Simplify  $\frac{32q^9r^4}{4q^3r}$ 

Simplify 
$$\frac{1}{4q^3r}$$

$$\frac{32 \times 9^9 \times r^4}{4 \times 9^3 \times r} = \frac{32}{4} \times \frac{9^9}{9^3} \times \frac{r^4}{r}$$

$$= 8 \times 9^6 \times r^3$$

Laws of Indias


$$\frac{x^a}{x^b} = x^{a-b}$$

1 2 correct terms


(Total for Question 1 is 5 marks)

2 (a) Find the lowest common multiple (LCM) of 40 and 56

Prime Factorisation



LCH = Product of all numbers in venn diagram (only counting those 40 56 in the intersection once)



LCm = 2×2×2×5×7
= 280
(2)

 $A = 2^3 \times 3 \times 5$ 

$$B = 2^2 \times 3 \times 5^2$$

(b) Write down the highest common factor (HCF) of A and B.




HCF= product of Shared prime factors



(Total for Question 2 is 3 marks)

3 The line L is shown on the grid.



Find an equation for L.

Equation of a straight line 
$$y = mx + c$$

gradient:  

$$M = \Delta y = \frac{6}{2} = 3$$
 (1)

$$y=3x+c \text{ (1)}$$
when  $x=0$   $y=-6$   $y=3x-6$ 

(Total for Question 3 is 3 marks)

4 Raya buys a van for £8500 plus VAT at 20%

Raya pays a deposit for the van.

She then pays the rest of the cost in 12 equal payments of £531.25 each month.

Find the ratio of the deposit Raya pays to the total of the 12 equal payments. Give your answer in its simplest form.

A) Total cost of van:

B) Total cost of payments:

() Deposit: Van cost - payment cost

= 10200 - 6375

= £ 3825 (I)

C : B

Deposit: Total of 12 payments

3825 : 6375 (1)

Simplify Ratio

3825

$$\times 3$$
  $\left(\begin{array}{c} \frac{5}{3} \\ \end{array}\right)_{x3}$ 

Whole number ratio

3:5

(Total for Question 4 is 5 marks)



(a) Complete the table of values for  $y = x^2 - x - 6$ 

х	-3	-2	-1	0	1	2	3
у	6	0	-4	- 6	-6	- 4	0


Substitute each & value into y=x2 to obtain the corresponding y value (same column in table)

e.g. when 
$$x = -2$$
  $y = x^2 - x - 6$   
=  $(-2)^2 - (-2) - 6$   
=  $4 + 2 - 6$   
 $y = 0$ 

(2)

(b) On the grid, draw the graph of  $y = x^2 - x - 6$  for values of x from -3 to 3

(2)



- (c) Use your graph to find estimates of the solutions to the equation  $x^2 x 6 = -2$
- · Draw the line y=-2 onto the graph (1)
- \* Find the x values of the 2 points at which the line y=-2 and the curve  $y=-\infty^2-\infty-6$  cross
- $\overline{y} = x^2 x 6$ -1.6 and 2.6

(Total for Question 5 is 6 marks)

A force of 70 newtons acts on an area of 20 cm<sup>2</sup>

The force is increased by 10 newtons. The area is increased by 10 cm<sup>2</sup>

$$pressure = \frac{force}{area}$$

Helen says,

"The pressure decreases by less than 20%"

Is Helen correct?
You must show he Initial Pressure:

You must show he'
$$P = \frac{F}{A} = \frac{70}{20} = 3.5$$
These values both have the Same unit (Ncm-1)
$$P = \frac{F}{A} = \frac{70 + 10}{20 + 10} = \frac{80}{30} = 2.6$$

$$P = \frac{F}{A} = \frac{70 + 10}{20 + 10} = \frac{80}{30} = 2.6$$

$$P = \frac{F}{A}$$

$$\frac{70 + 10}{20 + 10} = \frac{80}{30} = 2.6$$

$$\frac{80 + 10}{30} = \frac{80 + 10}{30}$$

New Pressure
$$P = \frac{F}{A} = 2.8 > 2.6$$

No, Helen is incorrect. The decrease is greater than 20%


20% less than the initial pressure = 80% of initial pressure 3.5 × 0.8 = 2.8 (1)

No, Helen is incorrect. The decrease is greater than 20%

(Total for Question 6 is 3 marks)

7 11 10 e.g. for the new Point  $x_2$ , we need to travel 3 of the distance 8 betweenthe 7 Centre and x2  $20 \text{ (i. (0.1)} \rightarrow (3.10)$ 6 is a shift of  $\binom{3}{9}$  $\binom{3}{9} \times \frac{1}{3} = \binom{3}{3}$  from 3 the centre point (0,1)

 $\therefore \propto_2 = (O+1, I+3)$  = (1, 4)



Enlarge shape A by scale factor  $\frac{1}{3}$  centre (0, 1)

Positive SF = expecting a Shape in the same orientation

SF < 1 = Smallershape - closer to centre

(Total for Question 7 is 2 marks)

60 people were asked if they prefer to go on holiday in Britain or in Spain or in Italy.

38 of the people were male.

11 of the 32 people who said Britain were female. 8 males said Italy.

12 people said Spain.

One of the females is chosen at random.

What is the probability that this female said Spain?

	B	5	I	tota	l
M			8	38	
F	11	3.3	ω	22	
Total	32	12	16	60	

Total for 
$$I = 60-32-12=16$$

F choosing 
$$I = 16 - 8 = 8$$
 (1)  
Total  $F = 60 - 38 = 22$ 

F choosing 
$$S = 22 - 11 - 8 = 3$$

P (the chosen female said spain) = 
$$\frac{\text{Females who chose spain}}{\text{Total number of females}} = \frac{3}{22}$$

(Total for Question 8 is 4 marks)

Jean invests £12000 in an account paying compound interest for 2 years.

In the first year the rate of interest is x%x . lo increase written as a decimal multiplier At the end of the first year the value of Jean's investment is £12336

In the second year the rate of interest is  $\frac{x}{2}$  %

What is the value of Jean's investment at the end of 2 years?

Set up equation for interest in year 1:
$$12000 \times \left(1 + \frac{\infty}{100}\right) = 12336$$

$$1 + \frac{\infty}{100} = 1.028...$$

$$\frac{\infty}{100} = 0.028...$$

$$\infty = 2.8$$

Find year 2 rate of interest:

$$\frac{\infty}{2} = \frac{2.8}{2} = 1.4 \%$$
1.4% increase written as a decimal multiplier

Year 2 interest:


$$12336 \times 1.014 = 12508.70$$
 1

'Compound interest' so we work with the Value at the end of the first year

£ 12508.70 🛈

(Total for Question 9 is 4 marks)

10 The vector **a** and the vector **b** are shown on the grid.



- (a) On the grid, draw and label vector -2a
- (b) Work out  $\mathbf{a} + 2\mathbf{b}$  as a column vector.
- $-2(\alpha) = -2\begin{pmatrix} 1\\2 \end{pmatrix} = \begin{pmatrix} -2 \times 1\\-2 \times 2 \end{pmatrix} = \begin{pmatrix} -2\\-4 \end{pmatrix}$

Vectors only show magnitude and direction, so can be drawn anywhere on the grid.

Work out a and b as column vectors:


$$a = \begin{pmatrix} 1 \\ 2 \end{pmatrix} \qquad b = \begin{pmatrix} 1 \\ -3 \end{pmatrix}$$

$$a + 2b = \begin{pmatrix} 1 \\ 2 \end{pmatrix} + 2 \begin{pmatrix} 1 \\ -3 \end{pmatrix} \qquad 0$$

$$= \begin{pmatrix} 1 \\ 2 \end{pmatrix} + \begin{pmatrix} 2 \times 1 \\ 2 \times -3 \end{pmatrix}$$

$$= \begin{pmatrix} 1 + 2 \\ 2 - 6 \end{pmatrix}$$

$$= \begin{pmatrix} 1+2\\2-6 \end{pmatrix}$$
$$= \begin{pmatrix} 3\\-4 \end{pmatrix}$$



(Total for Question 10 is 3 marks)

11 f and g are functions such that

$$f(x) = \frac{2}{x^2}$$
 and  $g(x) = 4x^3$ 

 $f(x) = \frac{2}{x^2} \quad \text{and} \quad g(x) = 4x^3$ (a) Find f(-5) Substitute x = -5 into f(x) function.

$$f(-5) = \frac{2}{(-5)^2} = \frac{2}{25}$$

(b) Find fg(1)

Composite function

'do g then do f' = 
$$f(g(x)) = f(g(1))$$


$$9(1) = 4 \times 1^{3} \boxed{1}$$

$$= 4$$


$$f(4) = \frac{2}{4^{2}} = \frac{2}{16} = \frac{1}{8}$$

(Total for Question 11 is 3 marks)

12



Graph B



Graph C

Otherwise go through origin

Graph D  $\begin{array}{c|c}
y & & \\
& & \\
O & & \\
\end{array}$ 

The graphs of y against x represent four different types of proportionality.

Match each type of proportionality in the table to the correct graph.

	onstant of proportionality	Type of proportionality	Graph letter
Straight _	$y = k \infty$ $y = m \infty + c$	$y \propto x$	В
line		$y \propto x^2$	D
	y=koc² + quadratic	$y \propto \sqrt{x}$	A
	+ quadratic	$y \propto \frac{1}{x}$	С

- 1) 2 correct
- (1) 4 correct


inversely proportional (Total for Question 12 is 2 marks)

as y increases, or decreases proportionally

— Will not go through the origin There are no real solutions for y when x=0  $\left(\frac{1}{0}X\right)$ , so the graph will tend to the yaxis.



13



- A, B, C and D are points on the circumference of a circle, centre O. FDE is a tangent to the circle.
- (a) Show that y-x=90You must give a reason for each stage of your working.

The tangent of the circle is perpendicular to its radius

$$\therefore \angle BDE = y = \infty + 90 \qquad y = \infty + 90$$

$$y - \infty = 90 \quad \text{as required (1)}$$
(3)

Dylan was asked to give some possible values for x and y.

He said,

"y could be 200 and x could be 110, because 200 - 110 = 90"

(b) Is Dylan correct?


You must give a reason for your answer.

No, y must be less than 180 asitis an angle in a triangle

(1)

(Total for Question 13 is 4 marks)

14 The distance-time graph shows information about part of a car journey.



Use the graph to estimate the speed of the car at time 5 seconds.

Finding the gradient of the tangent:  

$$M = Spee0 = \frac{\Delta y}{\Delta x} = \frac{68 - 4}{7.8 - 3.2}$$

$$= \frac{64}{4.6}$$


$$= 13.9$$

Any value in the range 11-19 m/s 13.9

(Total for Question 14 is 3 marks)

- 15 A darts team is going to play a match on Saturday and on Sunday. The probability that the team will win on Saturday is 0.45
  - If they win on Saturday, the probability that they will win on Sunday is 0.67 
    If they do **not** win on Saturday, the probability that they will win on Sunday is 0.35

(a) Complete the probability tree diagram.




(b) Find the probability that the team will win exactly one of the two matches.

Working across tree diagram, so we multiply 
$$P(win on Sat, (ose on sun) = 0.45 \times 0.33 = 0.1485 \bigcirc Probabilities.$$


0.341 (3)

(Total for Question 15 is 5 marks)



PhysicsAndMathsTutor.com

16 (a) On the grid, draw the graph of  $x^2 + y^2 = 12.25$  Circle 7 radius =  $\sqrt{12.25} = 3.5$ 



(b) Hence find estimates for the solutions of the simultaneous equations


$$x^2 + y^2 = 12.25$$
 $2x + y = 1$ 

Draw on grid - find where it Molts the circle

 $x = 2.0 \text{ y} = -2.9, \quad x = -1.2 \text{ y} = 3.3$ 
(3)

(Total for Question 16 is 5 marks)

17 The histogram shows information about the times taken by some students to finish a puzzle.



(a) Complete the frequency table for this information. Frequency = Frequency density X class width

	Time taken (t minutes)	Frequency	
	0 < <i>t</i> ≤ 5	4	
Class width_		4	0.4×10
= 15-5=10	$15 < t \leqslant 25$	6 (1)	0.6 × 10
	$25 < t \leqslant 30$	5	1 × 5
	$30 < t \leqslant 50$	4	6.2 × 20
			$\widehat{)} \qquad (2)$

(b) Find an estimate for the lower quartile of the times taken to finish the puzzle.

Time taken (t minutes)	Frequency	CF
0 < <i>t</i> ≤ 5	4	4
5 < <i>t</i> ≤ 15	4	8
$15 < t \leqslant 25$	6	14
$25 < t \leqslant 30$	5	19
$30 < t \leqslant 50$	4	23
•		∧

$$N = \xi f = 23$$

$$Q_1 = \frac{\Lambda+1}{4}$$
 to value

$$\frac{23+1}{4} = \frac{24}{4} = 6 \text{ th}$$

6th value is in the 5< t ≤ 15 interval

$$\frac{6-4}{4} = \frac{1}{2}$$

 $\frac{6-4}{4} = \frac{1}{2}$  halfway through interval

$$15-5=10$$
  $5+\left(\frac{10}{2}\right)=10$ 


lower bound

Ofinterval

minutes (2)

(Total for Question 17 is 4 marks)

**18** ABCDEFGH is a cuboid.




$$AB = 7.3 \text{ cm}$$
  
 $CH = 8.1 \text{ cm}$   
Angle  $BCA = 48^{\circ}$ 


Cuboid.

Find the size of the angle between *AH* and the plane *ABCD*. Give your answer correct to 1 decimal place.

Calculating Ac:



$$Sin \Theta = \frac{O}{h}$$
  
 $Sin 48 = \frac{7.3}{Ac}$   
 $Ac = \frac{7.3}{Sin 48} = 9.8231...$ 



tan 
$$O = \frac{O}{a}$$

tan  $O = \frac{81}{9.8231}$ .


 $O = \tan^{-1}\left(\frac{8.1}{9.8231...}\right)$ 

<u>()</u> 5


39.5


(Total for Question 18 is 4 marks)

19 Shape S is one quarter of a solid sphere, centre O.



Shape S







The volume of S is  $576\pi$  cm<sup>3</sup>

Find the surface area of S.

Give your answer correct to 3 significant figures.

You must show your working.

Finding 
$$\Gamma$$
:

 $4 \times 5767 = \frac{4}{3} \times 576$ 

SA of curved (outer) face:

SA of Other faces: 
$$\sim$$
 half circles with radius  $r$   $\frac{\pi r^2}{r^2} \times 2 = \pi r^2$ 

Total SA = 
$$\pi r^2 + \pi r^2 = 2\pi r^2$$

$$2\pi r^2 = 2 \times \pi \times 12^2$$
  
= 288 \pi  
= 90 4.779...

905 cm

(Total for Question 19 is 5 marks)

20 Martin did this question.

Rationalise the denominator of 
$$\frac{14}{2 + \sqrt{3}}$$

Here is how he answered the question.

$$\frac{14}{2+\sqrt{3}} = \frac{14 \times (2-\sqrt{3})}{(2+\sqrt{3})(2-\sqrt{3})}$$

$$= \frac{28-14\sqrt{3}}{4+2\sqrt{3}-2\sqrt{3}+3}$$

$$= \frac{28-14\sqrt{3}}{7}$$

$$= \frac{28-14\sqrt{3}}{7}$$

$$= 4-2\sqrt{3}$$

$$= 4-2\sqrt{3}$$

$$= \frac{14 \times (2-\sqrt{3})}{(2+\sqrt{3})}$$

$$= (2\times2)+(2\times\sqrt{3})+(2\times-\sqrt{3})+(\sqrt{3}\times-\sqrt{3})$$

$$= (2\times2)+(2\times\sqrt{3})+(2\times-\sqrt{3})+(\sqrt{3}\times-\sqrt{3})$$

$$= (3\times2)+(2\times\sqrt{3})+(2\times-\sqrt{3})+(2\times\sqrt{3})+(2\times-\sqrt{3})+(2\times$$

Martin's answer is wrong.

(a) Find Martin's mistake.

$$\sqrt{3} x - \sqrt{3} = -3 \text{ not } 3$$

(1)

Sian did this question.

Rationalise the denominator of  $\frac{5}{\sqrt{12}}$ 

Here is how she answered the question.

$$\frac{5}{\sqrt{12}} = \frac{5\sqrt{12}}{\sqrt{12} \times \sqrt{12}}$$

$$= \frac{5 \times 3\sqrt{2}}{12}$$

$$= \frac{5\sqrt{2}}{4}$$

$$\sqrt{12} = \sqrt{4}\sqrt{3}$$

$$= 2\sqrt{3}$$

Sian's answer is wrong.

(b) Find Sian's mistake.

$$\sqrt{12} = 2\sqrt{3}$$
, not  $3\sqrt{2}$ 

(1)

(Total for Question 20 is 2 marks)



21 Jackson is trying to find the density, in g/cm³, of a block of wood. The block of wood is in the shape of a cuboid.

He measures

the length as 13.2 cm, correct to the nearest mm the width as 16.0 cm, correct to the nearest mm the height as 21.7 cm, correct to the nearest mm

He measures the mass as 1970 g, correct to the nearest 5 g.

By considering bounds, work out the density of the wood. Give your answer to a suitable degree of accuracy.

You must show all your working and give a reason for your final answer.

$$m_{\nu} = 1972.5$$
  $m_{l} = 1967.5$   $\frac{5}{2} = 2.59$   $m_{ass} = 1970 \pm 2.59$ 

	3 sf	UB	LB
L	13.2	13.25	13.15
ω	16.0	16.05	15.95
h	21.7	21.75	21.65

$$V_{UB} = 13.25 \times 16.05 \times 21.75$$
  
= 4 625.4...

$$d = \frac{m}{v}$$

$$d_{v} = \frac{m_{v}}{v_{v}}$$

$$Q_{0} = \frac{1972.5}{4540.9...} = 0.4344...$$

$$Q_{0} = \frac{1967.5}{4625.4...} = 0.4254...$$

O. 43 g/cm<sup>3</sup>
Both the upper and lower bounds for density

(Total for Question 21 is 5 marks)

**TOTAL FOR PAPER IS 80 MARKS** 



**BLANK PAGE** 

