

Q1. A liquid flows continuously through a chamber that contains an electric heater. When the steady state is reached, the liquid leaving the chamber is at a higher temperature than the liquid entering the chamber. The difference in temperature is Δt .

Which of the following will increase Δt with no other change?

- A Increasing the volume flow rate of the liquid
- B Changing the liquid to one with a lower specific heat capacity
- C Using a heating element with a higher resistance
- D Changing the liquid to one that has a higher density

(Total 1 mark)

Q2. The temperature of a hot liquid in a container falls at a rate of 2 K per minute just before it begins to solidify. The temperature then remains steady for 20 minutes by which time all the liquid has all solidified.

What is the quantity $\frac{\text{Specific heat capacity of the liquid}}{\text{Specific latent heat of fusion}}$?

- A $\frac{1}{40} \text{ K}^{-1}$
- B $\frac{1}{10} \text{ K}^{-1}$
- C 10 K^{-1}
- D 40 K^{-1}

(Total 1 mark)

Q3. A fixed mass of gas occupies a volume V . The temperature of the gas increases so that the root mean square velocity of the gas molecules is doubled.
What will the new volume be if the pressure remains constant?

A $\frac{V}{2}$


B $\frac{V}{\sqrt{2}}$

C $2V$

D $4V$

(Total 1 mark)

Q4. **X** and **Y** are two gas bottles that are connected by a tube that has negligible volume compared with the volume of each bottle.



Initially the valve **W** is closed.

X has a volume $2V$ and contains hydrogen at a pressure of p .

Y has a volume V and contains hydrogen at a pressure of $2p$.

X and **Y** are both initially at the same temperature.

W is now opened. Assuming that there is no change in temperature, what is the new gas pressure?

A $\frac{2}{3}p$





B $\frac{5}{3}p$

C $\frac{4}{3}p$

D $\frac{3}{2}p$


(Total 1 mark)

Q5. Which one of the graphs below shows the relationship between the internal energy of an ideal gas (y -axis) and the absolute temperature of the gas (x -axis)?



(Total 1 mark)

Q6. The diagram shows the p - V diagram of an ideal hot-air engine. **WX** and **YZ** are isothermal changes.



Which line of the table below correctly indicates the nature of the work done **on** or **by** the air in each part of the cycle?

	WX	XY	YZ	ZW
A	zero	by	zero	on
B	by	zero	on	zero
C	zero	on	zero	by
D	on	zero	by	zero

(Total 1 mark)

Q7. The temperature of a room increases from 283K to 293K. The r.m.s. speed of the air molecules in the room increases by a factor of

- A** 1.02
- B** 1.04
- C** 1.41
- D** 2.00

(Total 1 mark)

Q8. A fixed mass of an ideal gas initially has a volume V and an absolute temperature T . Its initial pressure could be doubled by changing its volume and temperature to

- A $V/2$ and $4T$
- B $V/4$ and $T/2$
- C $2V$ and $T/4$
- D $4V$ and $2T$

(Total 1 mark)

Q9. A car of mass M travelling at speed V comes to rest using its brakes. Energy is dissipated in the brake discs of total mass m and specific heat capacity c . The rise in temperature of the brake discs can be estimated from

- A $\frac{mV^2}{2Mc}$
- B $\frac{2MV^2}{mc}$
- C $\frac{MV^2}{2mc}$
- D $\frac{2mc}{MV^2}$


(Total 1 mark)

Q10. Which one of the following is **not** an assumption about the properties of particles in the simple kinetic theory?

- A** $\frac{1}{2} m \langle v^2 \rangle$ is the average speed of the particles
- B** The forces between the particles are negligible except when particles collide
- C** The time spent by particles in collision is negligible compared with the time spent between collisions
- D** The volume of the particles is negligible compared to the volume of the container

(Total 1 mark)

Q11. The diagram shows a p - V graph for a fixed mass of gas. The volume increases from V_1 to V_2 while the pressure falls from p_1 to p_2 .




Which one of the paths **A**, **B**, **C** or **D** will result in the greatest amount of work being done by the gas?

(Total 1 mark)

Q12. In the diagram the dashed line **X** shows the variation of pressure, p , with absolute temperature, T , for 1 mol of an ideal gas in a container of fixed volume.

Which line, **A**, **B**, **C** or **D** shows the variation for 2 mol of the gas in the same container?




(Total 1 mark)

Q13. A raindrop of mass m falls to the ground at its terminal speed v . The specific heat capacity of water is c and the acceleration of free fall is g . Given that 25% of the energy is retained in the raindrop when it strikes the ground, what is the rise in temperature of the raindrop?

- A $\frac{mv^2}{8c}$
- B $\frac{v^2}{4mc}$
- C $\frac{mg}{4c}$
- D $\frac{v^2}{8c}$

(Total 1 mark)

Q14. The graph shows the relation between the product *pressure × volume*, pV , and temperature, θ , in degrees celsius for 1 mol of an ideal gas for which the molar gas constant is R .



Which one of the following expressions gives the gradient of this graph?

- A $\frac{1}{273}$
- B $\frac{pV}{\theta}$
- C $\frac{pV}{(\theta - 273)}$
- D R

(Total 1 mark)

Q15. At a certain temperature, the root-mean-square speed of the molecules of a fixed volume of an ideal gas is c . The temperature of the gas is changed so that the pressure is halved. The root-mean-square speed of the molecules becomes

- A $\frac{c}{4}$
- B $\frac{c}{2}$
- C $\frac{c}{\sqrt{2}}$
- D $2c$

(Total 1 mark)

Q16. A $1.0 \text{ k}\Omega$ resistor is thermally insulated and a potential difference of 6.0 V is applied to it for 2.0 minutes. The thermal capacity of the resistor is 9.0 J K^{-1} . The rise in temperature, in K , is

- A** 1.3×10^{-3}
- B** 8.0×10^{-3}
- C** 0.48
- D** 0.80

(Total 1 mark)