

M1.(a) (i) ultraviolet / UV / UV light / ultra(-)violet ✓

1

(ii) electron(in ground state) has moved / in to higher (energy) level / shell / orbital / state OR up level / shell / orbital / state ✓

Ignore reference to photons

1

(iii) (free) electrons collide with orbital electrons / mercury electrons / electrons in atom ✓

transferring energy ✓

Ignore any reference to photons

2

(iv) (mercury) atoms have discrete / fixed / specific energy levels ✓
when electrons change levels they lose an exact / fixed / specific / discrete / set amount of energy OR photons emitted with exact / fixed / specific / discrete / set amount of energy ✓
(leading to photons of) fixed / particular / certain / discrete / specific / unique frequencies ✓

Each mark independent

Don't accept characteristic for 3rd mark

3

(b) (i) (use of $\lambda = c / f$)
 $f = 3 \times 10^8 / (254 \times 10^{-9})$ ✓
 $f = 1.18 \times 10^{15}$ (Hz) ✓

AE penalty if give answer to 1 sig fig

2

(ii) (use of $E = hf$)
 $E = 6.63 \times 10^{-34} \times 1.18 \times 10^{15} = 7.82 \times 10^{-19} \text{J}$ ✓
 $E = 7.82 \times 10^{-19} / 1.6 \times 10^{-19}$ ✓ = 4.9 (4.875) eV

CE part (i)

Range 4.8 – 5.0 acceptable

2

- (c) coating absorbs photons / uv light ✓
and re-emits (photons) of low(er) energy / long(er) wavelength / low(er) frequency ✓

Ignore any description of mechanism

2

[13]

- M2.** (a) (i) when electrons/atoms are in their lowest/minimum energy (state) **or** most stable (state) they (are in their ground state) ✓

1

- (ii) in either case an electron receives (exactly the right amount of) energy ✓
excitation promotes an (orbital) electron to **a higher energy/up a level** ✓
ionisation occurs (when an electron receives enough energy) **to leave** the atom ✓

3

- (b) electrons occupy discrete energy levels ✓
and need to absorb an exact amount of/enough energy to move to a higher level ✓
photons need to have certain frequency to provide this energy **or** $e = hf$ ✓
energy required is the same for a particular atom or have different energy levels ✓
all energy of photon absorbed ✓
in 1 to 1 interaction or clear **a/the photon** and **an/the electrons** ✓

4

- (c) energy = $13.6 \times 1.60 \times 10^{-19} = 2.176 \times 10^{-18}$ (J) ✓
 $hf = 2.176 \times 10^{-18}$ ✓
 $f = 2.176 \times 10^{-18} \div 6.63 \times 10^{-34} = 3.28 \times 10^{15}$ Hz ✓ 3 sfs ✓

4

[12]

- M3.** (a) (i) an electron/atom is at a higher level than the ground state **(1)**
or electron jumped/moved up to another/higher level 1
- (ii) electrons (or electric current) flow through the tube **(1)**
and collide with orbiting/atomic electrons or mercury atoms **(1)**
raising the electrons to a higher level (in the mercury atoms) **(1)** 3
- (iii) photons emitted from mercury atoms are in the **ultra violet** (spectrum) **or** high energy photons **(1)**
these photons are absorbed by the powder **or** powder changes frequency/wavelength **(1)**
and the powder emits photons in the visible spectrum **(1)**
incident photons have a variety of different wavelengths **(1)** max 3
- (b) (i) (use of $E = hf$)
 $-0.26 \times 10^{-18} - 0.59 \times 10^{-18} \text{ (1)} = 6.63 \times 10^{-34} \times f \text{ (1)}$
 $f = 0.33 \times 10^{-18} / (6.63 \times 10^{-34}) = 5.0 \times 10^{14} \text{ (Hz) (1)}$ 3
- (ii) **one** arrow between $n = 3$ and $n = 2$ **(1)** in correct direction **(1)** 2

[12]

- M4.** (a) lowest energy state/level that the electron can occupy
or state in which electron needs most energy to be released

B1

1

or the level of an unexcited electron (not lowest orbit)

(b) (i) force = mv^2/r or $mr\omega^2$ and $v = r\omega$

B1

$$8.1 \times 10^{-8} = 9.1 \times 10^{-31} \times v^2 / 5.3 \times 10^{-11}$$

or ($v^2 =$) 4.72×10^{12} seen

B1

$$2.17 \times 10^6 \text{ (m s}^{-1}\text{)}$$

B1

(ii) $\lambda = h/mv$ or $6.6 \times 10^{-34} / 9.1 \times 10^{-31} \times 2.2 \times 10^6$

C1

7

$$3.3 \times 10^{-10} \text{ m}$$

A1

(iii) circumference = $2\pi 5.3 \times 10^{-11} = 3.3 \times 10^{-10} \text{ m}$

M1

1 (allow e.c.f. from (ii))

A1

(c) (i) $1.9(4) \times 10^{-18} \text{ J}$

B1

(ii) $5.6 \times 10^{-19} \text{ J}$ (e.c.f. 2.5×10^{-18} – their (i))

B1

- (iii) energy difference $E = 3 \times 10^{-19}$ J
(condone any difference)

C1

$$E = hc/\lambda \text{ or } E = hf \text{ and } c=f\lambda$$

or their $E = 6.6 \times 10^{-34} \times 3.0 \times 10^8/\lambda$

C1

$$6.6 \text{ or } 6.7 \times 10^{-7} \text{ m}$$

A1

5

[13]

M5.A

[1]