

M1.C

[1]

- M2.(a) force between two (point) charges is
proportional to product of charges ✓
inversely proportional to square of distance between the charges ✓
Mention of force is essential, otherwise no marks.
Condone "proportional to charges".
Do not allow "square of radius" when radius is undefined.
Award full credit for equation with all terms defined.

2

- (b) V is inversely proportional to r [or $V \propto (-)1/r$] ✓
(V has negative values) because charge is negative
[or because force is attractive on + charge placed near it
or because electric potential is + for + charge and - for - charge] ✓
potential is defined to be zero at infinity ✓
Allow $V \times r = \text{constant}$ for 1st mark.

max 2

- (c) (i) $Q(= 4\pi\epsilon_0 rV) = 4\pi\epsilon_0 \times 0.125 \times 2000$
OR $\text{gradient} = Q / 4\pi\epsilon_0 = 2000 / 8$ ✓
(for example, using any pair of values from graph) ✓
 $= 28 (27.8) (\pm 1) \text{ (nC)}$ ✓
(gives $Q = 28 (27.8) \pm 1 \text{ (nC)}$) ✓

2

- (ii) at $r = 0.20\text{m}$ $V = -1250\text{V}$ and at $r = 0.50\text{m}$ $V = -500\text{V}$
so pd $\Delta V = -500 - (-1250) = 750 \text{ (V)}$ ✓
work done $\Delta W (= Q\Delta V) = 60 \times 10^{-9} \times 750$
 $= 4.5(0) \times 10^{-5} \text{ (J)} (45 \mu\text{J})$ ✓

(final answer could be between 3.9 and 5.1×10^{-5})

Allow tolerance of $\pm 50\text{V}$ on graph readings.

[Alternative for 1st mark:

$$\Delta V = \frac{27.8 \times 10^{-9}}{4\pi\epsilon_0} \times \left(\frac{1}{0.2} - \frac{1}{0.5} \right) \text{ (or similar substitution using 60 nC)}$$

instead of 27.8 nC:
use of 60 nC gives $\Delta V = 1620V$]

2

(iii) $E \left(= \frac{Q}{4\pi\epsilon_0 r^2} \right) = \frac{27.8 \times 10^{-9}}{4\pi\epsilon_0 \times 0.40^2} \checkmark = 1600 \text{ (1560) (V m}^{-1}\text{)} \checkmark$

[or deduce $E = \frac{V}{r}$ by combining $E = \frac{Q}{4\pi\epsilon_0 r^2}$ with $V = \frac{Q}{4\pi\epsilon_0 r} \checkmark$

from graph $E = \frac{625 \pm 50}{0.40} = 1600 \text{ (1560} \pm 130\text{) (V m}^{-1}\text{)} \checkmark$]

Use of $Q = 30 \text{ nC}$ gives $1690 \text{ (V m}^{-1}\text{)}$.

Allow ecf from Q value in (i).

If $Q = 60 \text{ nC}$ is used here, no marks to be awarded.

2

[10]

- M3.(a)** (i) (Mass change in u) $1.71 \times 10^{-3} \text{ (u)}$
or (mass Be-7) – (mass He-3) – (mass He-4) seen with numbers

C1

$$2.84 \times 10^{-30} \text{ (kg)}$$

or Converts their mass to kg

Alternative 2nd mark:

Allow conversion of $1.71 \times 10^{-3} \text{ (u)}$ to MeV by multiplying by 931 (=1.59 (MeV)) **seen**

C1

Substitution in $E = mc^2$ condone their mass
difference in this sub but must have correct value for c^2
(3×10^8)² or 9×10^{16}

Alternative 3rd mark:

Allow their MeV converted to joules ($\times 1.6 \times 10^{-13}$) **seen**

C1

$$2.55 \times 10^{-13} \text{ (J) to } 2.6 \times 10^{-13} \text{ (J)}$$

Alternative 4th mark:

Allow 2.5×10^{-13} (J) for this method

A1
4

(ii) Use of $E=hc / \lambda$ **ecf**

C1

Correct substitution in rearranged equation with λ
subject **ecf**

C1

7.65×10^{-13} (m) to 7.8×10^{-13} (m) ecf

A1
3

(b) (i) Use of E_p formula:

C1

Correct charges for the nuclei **and** correct powers of 10

C1

$2.6(3) \times 10^{-13}$ J

A1
3

(ii) Uses $KE = 3 / 2 kT$: **or halves KE_T , $KE = 1.3 \times 10^{-13}$ (J)**
seen ecf

C1

Correct substitution of data **and** makes T subject **ecf**
Or uses KE_T value **and** divides T by 2

C1

6.35×10^9 (K) or 6.4×10^9 (K) or 6.28×10^9 (K) or 6.3×10^9 (K) **ecf**

A1
3

(c) (i) Deuteron / deuterium / hydrogen-2

B1

Triton / tritium / hydrogen-3

B1

2

(ii) Electrical heating / electrical discharge / inducing a current in plasma / use of e-m radiation / using radio waves (causing charged particles to resonate)

B1

1

[16]

M4.C

[1]

M5.D

[1]

M6.A

[1]

M7.B

[1]

M8.D

[1]

M9.D

[1]

M10.B

[1]

M11.C

[1]

M12. (a) work done [or energy needed] per unit charge [or (change in) electric pe per unit charge] ✓

on [or of] a (small) positive (test) charge ✓

in moving the charge from infinity (to the point) ✓

[not from the point to infinity] ✓

3

(b) (i) $V = \frac{Q}{4\pi\epsilon_0 r}$ gives $Q (= 4\pi\epsilon_0 rV) = 4\pi \times 8.85 \times 10^{-12} \times 0.30 \times 3.0$ ✓
 $= 1.0 \times 10^{-10}$ (C) ✓
to 2 sf only ✓

3

(ii) use of $V \propto \frac{1}{r}$ gives $V_m = \frac{V_L}{3}$ ✓ (= (+) 1.0 V)

1

$$(iii) \quad E \left(= \frac{Q}{4\pi\epsilon_0 r^2} \right) = \frac{1.0 \times 10^{-10}}{4\pi \times 8.85 \times 10^{-12} \times 0.60^2} \checkmark (= 2.50 \text{ V m}^{-1}) \quad 1$$

(c) (i) uniformly spaced vertical parallel lines which start and end on plates \checkmark
 relevant lines with arrow(s) pointing only downwards \checkmark 2

(ii) = 3.3(3) (V m⁻¹) \checkmark 1

(iii) part (b) is a radial field whilst part (c) is a uniform field \checkmark
 [or field lines become further apart between **L** and **M** but are equally spaced between **R** and **S**] 1

[12]